



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12Q 1/68	A1	(11) International Publication Number: WO 00/36142 (43) International Publication Date: 22 June 2000. (22.06.00)
(21) International Application Number: PCT/CA99/01177 (22) International Filing Date: 10 December 1999 (10.12.99) (30) Priority Data: 60/111,794 11 December 1998 (11.12.98) US (71) Applicant (for all designated States except US): VISIBLE GENETICS INC. [CA/CA]; 700 Bay Street, P.O. Box 333, Toronto, Ontario M5G 1Z6 (CA). (72) Inventor; and (75) Inventor/Applicant (for US only): SHIPMAN, Robert [CA/CA]; #2-6240 Montevideo Road, Mississauga, Ontario L5N 2N7 (CA). (74) Agent: DEETH WILLIAMS WALL; National Bank Building, Suite 400, 150 York Street, Toronto, Ontario M5H 3S5 (CA).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHOD AND KIT FOR THE CHARACTERIZATION OF ANTIBIOTIC-RESISTANCE MUTATIONS IN <i>MYCOBACTERIUM TUBERCULOSIS</i>		
(57) Abstract <p>Amplification and cycle sequencing primer sets have been developed for the detection and analysis of antibiotic resistance-associated mutations in defined regions of the <i>rpoB</i> (rifampin), <i>katG</i> (isoniazid), <i>oxyR-ahpC</i> PR (isoniazid), <i>mabA</i> (isoniazid), <i>rpsL/s12</i> (streptomycin), <i>16S/rrs</i> (streptomycin), <i>embB</i> (ethambutol), <i>pncA</i> (pyrazinamide), <i>gyrA</i> (ciprofloxacin) and 23S (azithromycin) genes of <i>Mycobacterium tuberculosis</i>. These primers can be used in a method for detection and characterization of <i>Mycobacterium tuberculosis</i> present in a sample. The method includes the steps of obtaining a sputum sample suspected of containing <i>M. tuberculosis</i>, performing a first sequencing procedure, with or without prior amplification, on the sample to detect the presence of <i>M. tuberculosis</i>, and if present to evaluate the <i>rpoB</i>, <i>katG</i>, <i>rpsL/s12</i> and 23S genes for the presence of antibiotic-resistance inducing mutations; and (c) if <i>M. tuberculosis</i> is detected in step (b), performing a second sequencing procedure, with or without prior amplification, on the sample to evaluate the additional genes for the presence of antibiotic-resistance inducing mutations.</p>		

